Detection of O-Linked-N-Acetylglucosamine Modification and Its Associated Enzymes in Human Degenerated Intervertebral Discs

Georgios Nikolaou¹, Aristeidis H. Zibis¹, Apostolos H. Fyllos¹, Antonios Katsioulis², Sotirios Sotiriou³, Anastasios Kotrotsios¹, Markos Sgantzios¹, Aikaterini Vassiou¹, Dimitrios L. Arvanitis¹

¹Department of Anatomy, Faculty of Medicine, University of Thessaly, Larissa, Greece

²Department of Hygiene and Epidemiology, Faculty of Medicine, University of Thessaly, Larissa, Greece

³Department of Embryology, Faculty of Medicine, University of Thessaly, Larissa, Greece

Study Design: Human herniated discs were obtained from discectomy specimens for the immunohistochemical detection of O-GlcNAc and O-GlcNAcase (OGA)/O-GlcNAc transferase (OGT).

Purpose: This study aimed to quantify the extent of O-GlcNAcylation and its associated enzymes (OGT/OGA) in human degenerated intervertebral discs.

Overview of Literature: The O-GlcNAcylation of nuclear, cytoplasmic, and mitochondrial proteins as well as the effects of such post-translational modifications are currently the focus of extensive research. O-GlcNAcylation is believed to contribute to the etiology of chronic illnesses by acting as a nutrient and stress sensor in the cellular environment. Mature intervertebral disc cells are chondrocyte-like cells, and O-GlcNAc has been shown to promote chondrocyte apoptosis *in vitro*. We believe that O-GlcNAcylation is a key regulator of disc degeneration.

Methods: Fifty-six specimens were fixed for 24 hours in a 10% solution of neutral-buffered formaldehyde, dehydrated, and embedded in paraffin. Tissue slices (4- μ m-thick) were used for hematoxylin-eosin staining and immunohistochemistry.

Results: We found that O-GlcNAcylation of cytoplasmic proteins was less than that of nuclear proteins in both single cells and cell clusters. Cytoplasmic O-GlcNAcylation occurs subsequent to nuclear O-GlcNAcylation and is directly proportional to disc degeneration. OGT and O-GlcNAc expression levels were identical in all specimens examined.

Conclusions: O-GlcNAc and OGA/OGT expression is shown to correlate for the first time with intervertebral disc cell degeneration. Increasing disc degeneration is associated with increasing O-GlcNAcylation in both nuclear and cytoplasmic proteins in human disc cells.

Keywords: Spine; Enzymes; Intervertebral disc degeneration; Immunohistochemistry

Introduction

The process of O-GlcNAcylation is defined as a post-trans-

lational, bidirectional, dynamic modification of serine and threonine residues within nuclear, mitochondrial, and cytoplasmic proteins by O-linked β -N-acetylglucosamine

Received Mar 6, 2017; Revised Apr 4, 2017; Accepted Apr 10, 2017

Corresponding author: Aristeidis Hristos Zibis

Department of Anatomy, Faculty of Medicine, University of Thessaly, Panepistimiou 3, Viopolis, Larissa 41334, Greece

Tel: +30-6977228250, Fax: +30-2410289266, E-mail: ahzibis@gmail.com

(O-GlcNAc). The O-GlcNAcylation substrate uridine diphospho-N-acetylglucosamine (UDP-GlcNAc) is directly produced by extracellular glucose metabolism via the hexosamine biosynthesis pathway, a major branch of the glycolytic pathway responsible for metabolizing 2%–5% of all cellular glucose [1]. As serum glucose levels rise in cases of diabetes or because of dietary uptake, more glucose is shunted into the hexosamine pathway, resulting in elevated levels of UDP-GlcNAc that can bind to proteins [2]. O-GlcNAcylation occurs post-translationally as a distinct form of glycosylation in both the nucleus and cytoplasm and is critical for cell transcription, proliferation, differentiation, and apoptosis. O-GlcNAcylation of serine and threonine residues is mediated by O-GlcNAc transferase (OGT) and reversed by O-GlcNAcase (OGA). The gene encoding OGT is highly conserved; its deletion is embryonically lethal [3,4]. O-GlcNAcylation is often suggested to serve as a nutrient sensor, as it is more common in hyperglycemic conditions, such as diabetes mellitus and metabolic syndrome [5,6].

The roles of biochemical factors in intervertebral disc degeneration and pain have received increased focus recently. The human intervertebral disc contains chondrocyte-like cells within the nucleus pulposus and inner annulus fibrosus, and fibroblast-like cells are found in the outer annulus fibrosus. These cells are responsible for the synthesis and regulation of the disc matrix and control of homeostasis between extracellular matrix synthesis and degradation. Increased protein O-GlcNAcylation plays an important role in the stress response [7]. Disc cells are capable of producing pro-inflammatory mediators, such as tumor necrosis factor α (TNF- α) and interleukin-1 (IL-1), which may directly stimulate discogenic back pain by promoting neoangiogenesis and sensitizing peripheral nociceptors. Further matrix degradation triggered by these agents may result in pain by impairing mechanical stability or structural integrity [8].

The purpose of this study was to investigate the O-GlcNAcylation of nuclear and cytoplasmic proteins in human degenerated intervertebral disc cells.

Materials and Methods

Fifty-six tissue samples were obtained from discectomy specimens of human herniated discs. The patient cohort included 30 men and 26 women, with a mean age of 45.6 years (range, 32 to 57 years). All patients presented with

extruded nucleus pulposus and symptoms of severe radiculopathy. Specimens were fixed for 24 hours in a 10% solution of neutral-buffered formaldehyde, dehydrated sequentially in 80%, 95%, and 100% solutions of ethanol and 100% xylene, and embedded in paraffin. Slices (4- μ m-thick) were used for hematoxylin-eosin staining and immunohistochemistry.

To identify O-GlcNAcylated proteins, we used the MAI-072 mouse monoclonal antibody RL2, which was obtained from Thermo Fisher Scientific (Waltham, MA, USA), at a 1:200 dilution, according to the manufacturer's instructions. To determine the localization of OGT, we used the PA5-13654 rabbit polyclonal antibody, which was obtained from Thermo Fisher Scientific, at a 1:100 dilution, according to the manufacturer's instructions. For the detection of OGA, we used the rabbit anti-MGEA5 antibody from Sigma-Aldrich Chemie GmbH (Taufkirchen, Germany). For immunohistochemistry experiments, the Genemed Kit Polymer HRP-containing goat antibody (South San Francisco, CA, USA) was used as a second antibody and for multicolor staining, according to the manufacturer's instructions.

Histological changes were graded according to the method described by Boos et al. [9]; we chose cell proliferation score (CPS) as a degeneration index. Immunohistochemistry grading was as follows: negative staining; weak staining (1+); moderate staining (2+); and intense staining (3+); for each intensity, if >75% of the cell population was stained, was considered high, if 25%–75% of the cell population was stained, it was considered moderate, and if <25% of the cell population was stained, it was considered low. All statistical analyses were performed using Microsoft Excel 2015 and SPSS ver. 21.0 (IBM Corp., Armonk, NY, USA).

The study was approved by the Institutional Review Board of University of Thessaly.

Results

The results of the immunohistochemistry experiments showed that there was nuclear and cytoplasmic staining of O-GlcNAc, OGT, and OGA in both single cells and cell clusters.

Staining results were identical for OGT and O-GlcNAcylated proteins. In all samples with a CPS of 5, nuclear staining was intense (3+) in >75% of single cells and cell clusters, whereas cytoplasmic staining was intense and

Fig. 1. (A) A large group of cells are shown. The overwhelming majority of cells shows intense nuclear and cytoplasmic staining ($\times 400$). (B) Single cells from an adjacent area of the same specimen show intense nuclear and cytoplasmic staining ($\times 400$).

Fig. 2. (A) A group of cells stained for O-GlcNAc transferase are shown. The great majority of cells shows intense nuclear and cytoplasmic staining. Seven cells remain unstained (blue nuclei) ($\times 400$). (B) Single cells from an adjacent area of the same specimen show intense nuclear and cytoplasmic staining ($\times 400$).

high ($>75\%$ of cells) in 46% of samples and moderate (2+) and high in the remaining samples. In all samples with a CPS of 4, nuclear staining was intense and high in single cells and cell clusters, **whereas cytoplasmic staining** was intense and high in only 16% of samples and moderate and high in the remaining samples. In samples with a CPS of 3, nuclear staining was intense and high in 77% of samples and moderate and high in the remaining four samples. Furthermore, in samples with a CPS of 3, cytoplasmic staining was moderate and high in 14 of 18 samples, and weak (1+) and high in the remaining four

Fig. 3. (A) Immunohistochemistry for O-GlcNAcase (OGA) is shown. A group of three cells with intense nuclear and weak cytoplasmic staining are shown ($\times 400$). (B) A group of herniated intervertebral disc cells are shown. Immunohistochemistry for OGA is shown. All cells show negative nuclear staining (blue nuclei). Few cells with weak cytoplasmic staining (brown color) are shown ($\times 400$).

samples.

Out of 26 samples with a CPS of 5, nuclear OGA staining was intense and high in 23% of samples, intense and moderate (25%–75% of cells) in 23% of samples, weak and low ($<25\%$ of cells) in four samples, and completely negative in the remaining 10 samples. Cytoplasmic staining was weak and high in six samples, weak and moderate in six samples, and negative in 14 samples. In samples with a CPS of 4, both nuclear and cytoplasmic staining was negative in all 12 samples. In samples with a CPS of 3, nuclear staining was intense and high in 11% of samples, intense and moderate in two samples, moderate and high in two samples, weak and low in two samples, and completely negative in the remaining 10 samples. Cytoplasmic staining was weak and high in 22% of samples, weak and moderate in two samples, and negative in the remaining 12 samples (Figs. 1–5).

Kendall's tau-b correlation coefficient was used to examine the relationship between the O-GlcNAc staining of nuclear and cytoplasmic proteins and CPS. A statistically significant moderate association was found between the O-GlcNAc staining of nuclear proteins and CPS (Kendall's coefficient=0.334, $p=0.027$), and a significantly positive strong association was found between the O-GlcNAc staining of cytoplasmic proteins and CPS (Kendall's coefficient=0.513, $p<0.001$). Moreover, a significantly positive association was found between the O-GlcNAc staining of nuclear and cytoplasmic proteins (Kendall's coeffi-

Fig. 4. Nuclear expression of O-GlcNAc/O-GlcNAc transferase (OGT) and O-GlcNAcase (OGA) is shown. CPS, cell proliferation score as defined in the text; H, high; M, moderate; L, low; N, none.

Fig. 5. Cytoplasmic expression of O-GlcNAc/O-GlcNAc transferase (OGT) and O-GlcNAcase (OGA) is shown. CPS, cell proliferation score as defined in the text; H, high; M, moderate; L, low; N, none.

cient=0.530, $p=0.025$) with a low agreement (25%).

Discussion

Intervertebral disc cells can be subjected to stress in the cellular environment due to avascularity and mechanical loading conditions. Such cells reside in a vulnerable environment characterized by hypoxia and low-glucose availability, as their nutrition is dependent upon diffusion from the neighboring capillary circulation [10,11]. Chondrocytes tend to proliferate and become hypertrophic under stress. Normal chondrocyte-like intervertebral disc cells exist in isolation, whereas degenerated disc cells are housed in stress-induced clusters that are derived from a

single progenitor cell [12]. Stress often stimulates cellular glucose uptake, and increased protein O-GlcNAcylation has been shown to play an important role in this stress response [7], either through OGA inhibition or OGT overexpression. The elevation of O-GlcNAc levels before, or immediately after, the induction of cellular injury is protective in models of heat stress, oxidative stress, endoplasmic reticulum stress, hypoxia, ischemia reperfusion injury, and trauma hemorrhage [13,14]. Conversely, a sustained increase in O-GlcNAcylation contributes to the development of vascular and cardiac dysfunction [15]. O-GlcNAc represents not only a nutrient sensor but also a stress sensor. Cells can consequently modify O-GlcNAc levels to both respond and adapt to the cellular environ-

ment. The purpose of the current study was to quantify O-GlcNAc in human degenerated disc cells.

O-GlcNAc was present in both the nucleus and cytoplasm of degenerated herniated intervertebral disc cells at different levels, according to the degree of disc degeneration, which is reflected by the CPS. O-GlcNAc has a predominantly nuclear distribution, is associated with chromatin, and is found to bind with many cytoplasmic and mitochondrial proteins [16-18]. In all samples with a CPS of 4 or 5, nuclear staining was both intense and high. There were also no differences in the staining of single cells and cell clusters as a result of stress-induced proliferation. These findings suggest that either O-GlcNAc or stress similarly affect both single cells and cell clusters, or immunohistochemistry experiments are unable to resolve such subtle cellular differences. Nuclear staining was more intense in samples with a higher CPS in a statistically significant manner. The fact that nuclear staining was reduced in samples with a CPS of 3 compared with those with a higher CPS suggests that increased disc degeneration correlates with greater O-GlcNAcylation of nuclear proteins.

O-GlcNAc staining of cytoplasmic proteins was less than that of nuclear proteins. Cytoplasmic staining was intense and high in only 46% and 16% of samples with a CPS of 5 and 4, respectively. However, nuclear staining was intense in every sample with a CPS of 5 or 4. No sample with a CPS of 3 had intense cytoplasmic staining, whereas 77% of such samples had intense nuclear staining. Furthermore, in samples with a CPS of 3, there was weak cytoplasmic staining in 23% of samples, compared with samples with a CPS of 5 or 4. Therefore, cytoplasmic O-GlcNAcylation occurs subsequent to nuclear O-GlcNAcylation and is directly proportional to disc degeneration.

OGT and O-GlcNAc expression levels were identical in all specimens examined. Identical increases in nuclear staining of these proteins were observed as intervertebral disc degeneration increased, with staining of cytoplasmic proteins less than that of nuclear proteins. However, a different pattern was observed for OGA expression. Negative OGA expression was observed in samples with a CPS of 4, whereas samples with a CPS of 3 or 5 displayed a range of OGA staining results. OGA cytoplasmic expression occurred subsequent to its nuclear expression. Therefore, OGA expression cannot be characterized as the opposite of O-GlcNAc or OGT expression, considering its negative

expression in all samples with a CPS of 4.

An example of an inflammatory role for a specific O-GlcNAcylated protein is p65, a subunit of nuclear factor κ B (NF- κ B), in rheumatoid arthritis (RA). Its effects were studied in both fibroblast-like synoviocytes (FLS) and a mouse model of collagen-induced arthritis (CIA). Cells were first treated with an OGA inhibitor and then with TNF- α , resulting in the enhancement of O-GlcNAcylation. It was shown that hyper-O-GlcNAcylation increased both cell proliferation and expression levels of pro-inflammatory cytokines associated with RA *in vitro* and aggravated the severity of CIA *in vivo*. In contrast, hypo-O-GlcNAcylation in response to OGA overexpression significantly attenuated inflammation in TNF- α -stimulated FLS. O-GlcNAcylation of p65 enhanced inflammatory processes in TNF- α -stimulated FLS via increases in the nuclear translocation of p65, protein-DNA binding, and transcriptional activity, which increased the expression levels of downstream genes [19]. In contrast, another study reported that glucosamine improved cardiac function in rats following hemorrhagic shock and attenuated the activation of NF- κ B, which plays a central role in regulating the release of many cytokines, including TNF- α [20]. Finally, elevated O-GlcNAcylation appeared to significantly enhance colitis and colitis-associated colon cancer by modulating the DNA-binding activity of p65 *in vivo* [21].

It appears that hyperglycemia (in conditions of diabetes or dietary excess) has detrimental effects on human chondrocytes. Mature intervertebral disc cells are chondrocyte-like cells [22] that undergo the typical cascade of chondrogenic differentiation events (cell proliferation, extracellular matrix synthesis, cellular hypertrophy, matrix mineralization, vascular invasion, and eventually apoptosis), which allows cartilage remodeling into bone [23]. Diabetes can cause intervertebral disc degeneration by accelerating NP cell senescence and apoptosis [24]. Additionally, O-GlcNAc accumulation has been shown to increase chondrocyte apoptosis *in vitro* [25]. Furthermore, levels of O-GlcNAcylated proteins were found to increase four-fold in human osteoarthritic (OA) cartilage compared with healthy cartilage, a condition partly induced by IL-1 [26]. The increase in O-GlcNAcylation observed in OA cartilage was associated with a redistribution of expressed OGT isoforms, and an increase in OGA expression was also observed. Increased expression levels of OGT have been associated with O-GlcNAc accumulation

in response to both acute stress and chronic damage [27]. In human OA chondrocytes, O-GlcNAc accumulation parallels increases in both OGT and OGA, suggesting that OGT and OGA levels may fluctuate in response to modifications of O-GlcNAc levels [25], similar to our results.

In recent research on diabetic rats, the apoptotic index of nucleus pulposus notochordal cells and the extent of transition of notochordal nucleus pulposus into pro-cartilaginous nucleus pulposus were significantly increased, confirming the hypothesis that hyperglycemia-induced O-GlcNAc expression promotes and accelerates chondrocyte apoptosis, resulting in degeneration. Expression levels of matrix metalloproteinases-1, -2, -3, and -13 were also significantly increased in diabetic rats [28]. A higher incidence of degenerative disc disease in patients with diabetes and at a younger age than nondiabetic populations has been reported [29], suggesting that hyperglycemia contributes to the accelerated degeneration of intervertebral disc cells via O-GlcNAc.

This study's major strength is the use of immunohistochemistry to detect O-GlcNAc in degenerated intervertebral disc cells from human noncadaveric tissues, which has not been previously published. The major limitations of this study are the lack of similar immunohistochemistry data from normal discs and a lack of correlation between magnetic resonance imaging results for disc degeneration and our immunohistochemistry data.

Conclusions

O-GlcNAcylated proteins mediate numerous cellular processes. Research on the role of O-GlcNAc in intervertebral disc degeneration could elucidate the mechanism of intervertebral disc degeneration and lead to its prevention in various metabolic states.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

ORCID

Aristeidis Hristos Zibis:

<https://orcid.org/0000-0001-7122-4317>

Apostolos Haralampos Fyllos:

<https://orcid.org/0000-0003-1384-2074>

Antonios Katsioulis:

<https://orcid.org/0000-0002-9337-7223>

Markos Sgantzos: <https://orcid.org/0000-0001-6337-4534>

Dimitrios Leonidas Arvanitis:

<https://orcid.org/0000-0001-6435-7461>

References

1. Chatham JC, Not LG, Fulop N, Marchase RB. Hexosamine biosynthesis and protein O-glycosylation: the first line of defense against stress, ischemia, and trauma. *Shock* 2008;29:431-40.
2. Ma J, Hart GW. Protein O-GlcNAcylation in diabetes and diabetic complications. *Expert Rev Proteomics* 2013;10:365-80.
3. Lubas WA, Frank DW, Krause M, Hanover JA. O-Linked GlcNAc transferase is a conserved nucleocytoplasmic protein containing tetratricopeptide repeats. *J Biol Chem* 1997;272:9316-24.
4. Shafi R, Iyer SP, Ellies LG, et al. The O-GlcNAc transferase gene resides on the X chromosome and is essential for embryonic stem cell viability and mouse ontogeny. *Proc Natl Acad Sci U S A* 2000;97:5735-9.
5. Hart GW, Housley MP, Slawson C. Cycling of O-linked beta-N-acetylglucosamine on nucleocytoplasmic proteins. *Nature* 2007;446:1017-22.
6. Olivier-Van Stichelen S, Hanover JA. You are what you eat: O-linked N-acetylglucosamine in disease, development and epigenetics. *Curr Opin Clin Nutr Metab Care* 2015;18:339-45.
7. Zachara NE, Hart GW. O-GlcNAc a sensor of cellular state: the role of nucleocytoplasmic glycosylation in modulating cellular function in response to nutrition and stress. *Biochim Biophys Acta* 2004;1673:13-28.
8. Risbud MV, Shapiro IM. Role of cytokines in intervertebral disc degeneration: pain and disc content. *Nat Rev Rheumatol* 2014;10:44-56.
9. Boos N, Weissbach S, Rohrbach H, Weiler C, Spratt KF, Nerlich AG. Classification of age-related changes in lumbar intervertebral discs: 2002 Volvo Award in basic science. *Spine (Phila Pa 1976)* 2002;27:2631-44.
10. Rajasekaran S, Babu JN, Arun R, Armstrong BR, Shetty AP, Murugan S. ISSLS prize winner: A study of diffusion in human lumbar discs: a serial magnetic resonance imaging study documenting the influence of the endplate on diffusion in normal and degenerate discs. *Spine (Phila Pa 1976)* 2004;29:2654-67.

11. Grunhagen T, Shirazi-Adl A, Fairbank JC, Urban JP. Intervertebral disk nutrition: a review of factors influencing concentrations of nutrients and metabolites. *Orthop Clin North Am* 2011;42:465-77.
12. Sharp CA, Roberts S, Evans H, Brown SJ. Disc cell clusters in pathological human intervertebral discs are associated with increased stress protein immunostaining. *Eur Spine J* 2009;18:1587-94.
13. Zachara NE, Molina H, Wong KY, Pandey A, Hart GW. The dynamic stress-induced "O-GlcNAc-ome" highlights functions for O-GlcNAc in regulating DNA damage/repair and other cellular pathways. *Amino Acids* 2011;40:793-808.
14. Arvanitis LD, Vassiou K, Kotrotsios A, Sgantzios MN. Hypoxia upregulates the expression of the O-linked N-acetylglucosamine containing epitope H in human ependymal cells. *Pathol Res Pract* 2011;207:91-6.
15. Dassanayaka S, Jones SP. O-GlcNAc and the cardiovascular system. *Pharmacol Ther* 2014;142:62-71.
16. Kelly WG, Hart GW. Glycosylation of chromosomal proteins: localization of O-linked N-acetylglucosamine in *Drosophila* chromatin. *Cell* 1989;57:243-51.
17. Love DC, Kochan J, Cathey RL, Shin SH, Hanover JA. Mitochondrial and nucleocytoplasmic targeting of O-linked GlcNAc transferase. *J Cell Sci* 2003;116:647-54.
18. Arvanitis DL, Arvanitis LD, Panourias IG, Kitsoulis P, Kanavros P. Mitochondria-rich normal, metaplastic, and neoplastic cells show overexpression of the epitope H recognized by the monoclonal antibody H. *Pathol Res Pract* 2005;201:319-24.
19. Kim HB, Lee SW, Mun CH, et al. O-linked N-acetylglucosamine glycosylation of p65 aggravated the inflammation in both fibroblast-like synoviocytes stimulated by tumor necrosis factor- α and mice with collagen induced arthritis. *Arthritis Res Ther* 2015;17:248.
20. Zou L, Yang S, Champattanachai V, et al. Glucosamine improves cardiac function following trauma-hemorrhage by increased protein O-GlcNAcylation and attenuation of NF- κ B signaling. *Am J Physiol Heart Circ Physiol* 2009;296:H515-23.
21. Yang YR, Kim DH, Seo YK, et al. Elevated O-GlcNAcylation promotes colonic inflammation and tumorigenesis by modulating NF- κ B signaling. *Oncotarget* 2015;6:12529-42.
22. Trout JJ, Buckwalter JA, Moore KC. Ultrastructure of the human intervertebral disc: II. Cells of the nucleus pulposus. *Anat Rec* 1982;204:307-14.
23. Kronenberg HM. Developmental regulation of the growth plate. *Nature* 2003;423:332-6.
24. Jiang L, Zhang X, Zheng X, et al. Apoptosis, senescence, and autophagy in rat nucleus pulposus cells: implications for diabetic intervertebral disc degeneration. *J Orthop Res* 2013;31:692-702.
25. Andres-Bergos J, Tardio L, Larranaga-Vera A, Gomez R, Herrero-Beaumont G, Largo R. The increase in O-linked N-acetylglucosamine protein modification stimulates chondrogenic differentiation both in vitro and in vivo. *J Biol Chem* 2012;287:33615-28.
26. Tardio L, Andres-Bergos J, Zachara NE, et al. O-linked N-acetylglucosamine (O-GlcNAc) protein modification is increased in the cartilage of patients with knee osteoarthritis. *Osteoarthritis Cartilage* 2014;22:259-63.
27. Zachara NE. The roles of O-linked beta-N-acetylglucosamine in cardiovascular physiology and disease. *Am J Physiol Heart Circ Physiol* 2012;302:H1905-18.
28. Won HY, Park JB, Park EY, Riew KD. Effect of hyperglycemia on apoptosis of notochordal cells and intervertebral disc degeneration in diabetic rats. *J Neurosurg Spine* 2009;11:741-8.
29. Sakellariadis N. The influence of diabetes mellitus on lumbar intervertebral disk herniation. *Surg Neurol* 2006;66:152-4.