The Time Course Changes in Bone Metabolic Markers after Administering the Anti-Receptor Activator of Nuclear Factor-Kappa B Ligand Antibody and Drug Compliance among Patients with Osteoporosis

Kazuhide Inage¹, Sumihisa Orita¹, Kazuyo Yamauchi¹, Yoshihiro Sakuma²,
Go Kubota¹, Yasuhiro Oikawa³, Takeshi Sainoh¹, Jun Sato¹, Kazuki Fujimoto¹,
Yasuhiro Shiga¹, Kazuhisa Takahashi¹, Seiji Ohtori¹

¹Department of Orthopedic Surgery, Graduate School of Medicine, Chiba University, Chiba, Japan

²Department of Orthopedic Surgery, National Hospital Organization Chiba Medical Center, Chiba, Japan

³Department of Orthopedic Surgery, Teikyo University Chiba Medical Center, Chiba, Japan

Asian Spine J 2015;9(3):338-343

<http://dx.doi.org/10.4184/asj.2015.9.3.338>

Corrections for Fig. 2, Fig. 3, and Fig. 4 in page 340 are needed.

There were some mistakes in the numerical values of the graphs. We apologize for any inconvenience that this may have caused.

• Before correction

Fig. 2. P1NP levels before and 1 month after administering the anti-RANKL antibody. P1NP, N-terminal propeptide of type 1 collagen; RANKL, receptor activator of nuclear factor-kappa B ligand; NS, not significant.

• After correction

Fig. 2. P1NP levels before and 1 month after administering the anti-RANKL antibody. P1NP, N-terminal propeptide of type 1 collagen; RANKL, receptor activator of nuclear factor-kappa B ligand; NS, not significant.

• Before correction

Fig. 3. The averages YAM level for the anti-RANKL antibody before and 6 months after treatment. YAM, young adult mean; RANKL, receptor activator of nuclear factor-kappa B ligand.

• After correction

Fig. 3. The averages YAM level for the anti-RANKL antibody before and 6 months after treatment. YAM, young adult mean; RANKL, receptor activator of nuclear factor-kappa B ligand.

• Before correction

Fig. 4. The time course changes by using the back pain VAS before treatment and 1, 2, 3, and 6 months after treatment. VAS, visual analogue scale; NS, not significant.

• After correction

Fig. 4. The time course changes by using the back pain VAS before treatment and 1, 2, 3, and 6 months after treatment. VAS, visual analogue scale; NS, not significant.